


Sucrose

Add the following:

- Portions of the monograph text that are national *USP* text, and are not part of the harmonized text, are marked with symbols (♦) to specify this fact. 1S (NF32)



$C_{12}H_{22}O_{11}$ 342.30
 α -D-Glucopyranoside, β -D-fructofuranosyl-sucrose [57-50-1].

DEFINITION

Sucrose is a sugar obtained from *Saccharum officinarum* Linné (Fam. Gramineae), *Beta vulgaris* Linné (Fam. Chenopodiaceae), and other sources. It contains no added substances.

IMPURITIES

Delete the following:

- RESIDUE ON IGNITION (281)**

Sample: 5 g

Acceptance criteria: NMT 0.05% 1S (NF32)

Delete the following:

- CHLORIDE AND SULFATE, Chloride (221)**

Sample: 2.0 g

Acceptance criteria: 0.0035%; the *Sample* shows no more chloride than corresponds to 0.10 mL of 0.020 N hydrochloric acid. 1S (NF32)

Delete the following:

- CHLORIDE AND SULFATE, Sulfate (221)**

Sample: 5.0 g

Acceptance criteria: 0.006%; the *Sample* shows no more sulfate than corresponds to 0.30 mL of 0.020 N sulfuric acid. 1S (NF32)

Delete the following:

- CALCIUM**

Analysis: To 10 mL of a solution (1 in 10) add 1 mL of ammonium oxalate TS.

Acceptance criteria: The solution remains clear for at least 1 min. 1S (NF32)

Delete the following:

- HEAVY METALS (231)**

Sample: 4.0 g

Analysis: Dissolve the *Sample* in 15 mL of water, add 1 mL of 0.12 N hydrochloric acid, and dilute with water to 25 mL.

Acceptance criteria: NMT 5 ppm 1S (NF32)

Add the following:

- SULFITE**

Sample solution: 400 mg/mL of Sucrose in freshly prepared distilled water

Sulfite standard solution (80 ppm SO_2): 0.1575 mg/mL of anhydrous sodium sulfite in freshly prepared distilled water

Reference solution: Dissolve 4.0 g of Sucrose in freshly prepared distilled water, add 0.5 mL of *Sulfite standard solution* (80 ppm SO_2), and dilute with freshly prepared distilled water to 10.0 mL.

Blank: Freshly prepared distilled water

Analysis: Determine the sulfite content by a suitable enzymatic method based on the following reactions. Sulfite is oxidized by sulfite oxidase to sulfate and hydrogen peroxide, which in turn is reduced by nicotinamide-adenine dinucleotide-peroxidase in the presence of reduced nicotinamide-adenine dinucleotide (NADH). The amount of NADH oxidized is proportional to the amount of sulfite. Separately introduce 2.0 mL each of the *Sample solution*, *Reference solution*, and *Blank* in 10-mm cuvettes, and add the reagents as described in the kit instructions.¹ Measure the absorbance at the maximum at about 340 nm before and at the end of the reaction time, and subtract the value obtained with the *Blank*.

Acceptance criteria: The absorbance difference of the *Sample solution* is NMT half the absorbance difference of the *Reference solution*. 1S (NF32)

SPECIFIC TESTS

Add the following:

- APPEARANCE OF SOLUTION**

Sample solution: 500 mg/mL of Sucrose in water.

[NOTE—Set a portion of this solution aside for the tests for *Dextrins* and *Reducing Sugars*.]

Hydrazine sulfate solution: 10 mg/mL of hydrazine sulfate in water. Allow to stand for 4–6 h.

Hexamethylenetetramine solution: In a 100-mL ground glass-stoppered flask dissolve 2.5 g of hexamethylenetetramine in 25.0 mL of water.

Primary opalescent suspension: To the *Hexamethylenetetramine solution* in the flask add 25.0 mL of *Hydrazine sulfate solution*. Mix, and allow to stand for 24 h. This suspension is stable for 2 months, provided it is stored in a glass container free from surface defects. The suspension must not adhere to the glass and must be well mixed before use.

Standard of opalescence: *Primary opalescent suspension* in water (3 in 200). This suspension is freshly prepared and may be stored for up to 24 h.

Reference suspension I: *Standard of opalescence* and water (5:95)

Acceptance criteria: The clarity of the *Sample solution* is the same as that of water or its opalescence is not more pronounced than that of *Reference suspension I*.

1S (NF32)

¹ Test kit for sulfite determination may be ordered from Boehringer Mannheim Roche/R-Biopharm, Catalog #10 725 854 035.

2 Sucrose

Add the following:

• CONDUCTIVITY

Sample solution: 313 mg/mL of Sucrose in freshly boiled and cooled water

Apparatus: Use a conductivity meter or resistivity meter that measures the resistance of the column of liquid between the electrodes of the immersed measuring device. The apparatus is supplied with alternating current to avoid the effects of electrode polarization. It is equipped with a temperature compensation device or a precision thermometer.

Calibration: Choose a conductivity cell that is appropriate for the conductivity of the solution to be examined. The higher the expected conductivity, the higher the cell constant that must be chosen so that the value measured, R , is as large as possible for the apparatus used. Commonly used conductivity cells have cell constants on the order of 0.1 cm^{-1} , 1 cm^{-1} , and 10 cm^{-1} . Use a standard solution of potassium chloride that is appropriate for the measurement. Rinse the cell several times with water that has been previously boiled and cooled to room temperature and at least twice with the potassium chloride solution used for the determination of the cell constant of the conductivity cell. Measure the resistance of the conductivity cell using the potassium chloride solution at $20 \pm 0.1^\circ$.

The constant, C (in cm^{-1}), of the conductivity cell is given by the expression:

$$C = R_{KCl} \times K_{KCl}$$

R_{KCl} = measured resistance ($M\Omega$)

K_{KCl} = conductivity of the standard solution of potassium chloride used ($\mu\text{S} \cdot \text{cm}^{-1}$)

The measured constant, C , of the conductivity cell must be within 5% of the given value.

Analysis

Sample: *Sample solution*

Measure the conductivity of the *Sample solution* (C_1), while gently stirring with a magnetic stirrer, and that of the water used for preparing the *Sample solution* (C_2). The readings must be stable within 1% over a period of 30 s.

Calculate the conductivity of the *Sample solution* from the expression:

$$\text{Result} = C_1 - (0.35 \times C_2)$$

C_1 = conductivity of the *Sample solution*

C_2 = water used for preparing the *Sample solution*

Acceptance criteria: NMT $35 \mu\text{S} \cdot \text{cm}^{-1}$ at 20° ■1S (NF32)

Change to read:

• OPTICAL ROTATION, *Specific Rotation* (781S)

Sample solution: 260 mg/mL (ERR 1-May-2012)

Acceptance criteria: +66.3 to +67.0 at 20° (ERR 1-May-2012)

Add the following:

• COLOR VALUE

Sample solution: Dissolve 50.0 g in 50.0 mL of water. Mix, filter (diameter of pores, $0.45 \mu\text{m}$), and degas.

Analysis: Measure the absorbance at 420 nm, using a cell of at least 4 cm (a cell length of 10 cm or more is preferred).

Calculate the *Color Value* using the expression:

$$\text{Result} = (A \times 1000)/(b \times c)$$

A = absorbance measured at 420 nm

b = cell path length (cm)

c = concentration of the solution (g/mL), calculated from the refractive index of the solution. Use *Table 1*, and interpolate the values if necessary. The absolute difference between two results is NMT 3.

Table 1

n_D^{20}	c (g/mL)
1.4138	0.570
1.4159	0.585
1.4179	0.600
1.4200	0.615
1.4221	0.630
1.4243	0.645
1.4264	0.661

Acceptance criteria: NMT 45 if labeled as parenteral grade; NMT 75 for nonparenteral grade ■1S (NF32)

Add the following:

• DEXTRINS

[NOTE—If intended for use in the preparation of large-volume infusions, it complies with the test for *Dextrins*.]

Sample solution: Prepare as directed in the test for *Appearance of Solution*.

Analysis: To 2 mL of the *Sample solution* add 8 mL of water, 0.05 mL of dilute hydrochloric acid (73 g/L of HCl), and 0.05 mL of 0.05 M iodine.

Acceptance criteria: The solution remains yellow.

■1S (NF32)

Add the following:

• REDUCING SUGARS

Sample solution: Prepare as directed in the test for *Appearance of Solution*.

Analysis: To 5 mL of the *Sample solution* in a test tube, about 150-mm long and 16-mm in diameter, add 5 mL of water, 1.0 mL of 1 M sodium hydroxide, and 1.0 mL of a 1-g/L solution of methylene blue. Mix, and place in a water bath. After exactly 2 min, take the tube out of the bath, and examine the solution immediately.

Acceptance criteria: The blue color does not disappear completely, ignoring any blue color at the air/solution interface. ■1S (NF32)

Add the following:

• LOSS ON DRYING (731)

Sample: 2.000 g

Analysis: Dry the *Sample* at 105° for 3 h.

Acceptance criteria: NMT 0.1% ■1S (NF32)

Add the following:

• BACTERIAL ENDOTOXINS TEST (85): Less than 0.25 IU/mg

[NOTE—If intended for use in the preparation of large-volume infusions, it complies with the test for *Bacterial Endotoxins*.] ■1S (NF32)

Delete the following:

■ **INVERT SUGAR**

Sample solution: 200 mg/mL of Sucrose in water
[NOTE—Filter if necessary.]

Analysis: Place 50 mL of the clear liquid in a 250-mL beaker, add 50 mL of alkaline cupric tartrate TS, cover the beaker with a watch glass, and heat the mixture at such a rate that it comes to a boil in approximately 4 min, and boil for 2 min, accurately timed. Add at once 100 mL of cold, recently boiled water, and immediately collect the precipitated cuprous oxide on a tared filtering crucible containing a sintered-glass disk of medium pore size, or suitable equivalent. Thoroughly wash the residue on the filter with hot water, then with 10 mL of alcohol, and finally with 10 mL of ether, and dry at 105° for 1 h.

Acceptance criteria: The weight of the cuprous oxide does not exceed 112 mg. ■^{1S} (NF32)

ADDITIONAL REQUIREMENTS

- ◆ **PACKAGING AND STORAGE:** Preserve in well-closed containers. ◆

Add the following:

- **LABELING:** The label states, where applicable, that the substance is suitable for use in the manufacture of large-volume parenteral dosage forms. ■^{1S} (NF32)